



ISSN: 2091-2749 (Print)  
2091-2757 (Online)

#### Correspondence

Ms. Sulochana Basukala  
Ward In-charge  
Patan Hospital  
Patan Academy of Health  
Sciences, Lalitpur, Nepal  
Email:  
sulochanabasukala@gmail.com

#### Peer Reviewers

Lect. Dr. Pawan Sharma  
Patan Academy of Health  
Sciences

Prof. Dr. Jay N Shah  
Patan Academy of Health  
Sciences

#### Submitted

20 Apr 2019

#### Accepted

05 Jun 2019

#### How to cite this article

Sulochana Basukala. Care giver burden among spouse of psychiatric patients attending psychiatry outpatient department of a teaching hospital. Journal of Patan Academy of Health Sciences. 2019Jun;6(1):72-77.

## Care giver burden among spouse of psychiatric patients attending psychiatry outpatient department of a teaching hospital

Sulochana Basukala\*

Ward In-charge, Patan Hospital, Patan Academy of Health Sciences, Lalitpur, Nepal

*\*During study period, author was Masters of Nursing student at Lalitpur Nursing Campus, School of Nursing and Midwifery, Patan Academy of Health Sciences, Lalitpur, Nepal.*

### Abstract

**Introductions:** Caregiver burden in mental illness is a global issue. Burden is more significant and high among spouse than other caregivers. Spouse face illness-specific burdens of feeling powerlessness, fear, marital stress and partnership and family roles.

**Methods:** A cross-sectional study was conducted in outpatient department of psychiatry in Patan hospital, Patan Academy of Health Sciences, Nepal, from 26<sup>th</sup> August to 23<sup>rd</sup> September, 2018. Non-probability purposive sampling technique was used. Data was collected from spouses of psychiatric patients by face-to-face interview using Burden Assessment Schedule (BAS) to assess the caregiver burden of spouses of psychiatric patients.

**Results:** Out of 160 spouses interviewed, 94 (59%) had severe burden, 13 (9%) very severe, and 53 (34%) moderate. Burden was significantly associated with respondents' relation to patient, educational status, income and diagnosis. The respondent's age, occupation, duration of caring, and place of residence had no significant association with burden.

**Conclusions:** More than half of the spouses of psychiatric patients had severe care giver burden.

**Keywords:** burden, psychiatric patient, spouse

## Introductions

Mental illnesses are rising worldwide.<sup>1</sup> It is a top cause of global burden of disease.<sup>2</sup> It affects individuals as well as their caregivers.<sup>3</sup> The burden of care, usually falls on the spouse in marital relationships.<sup>4</sup> Living with and caring for psychiatric spouse is an enduring stressor and causes considerable burdens.<sup>5</sup> Caregiver burden refers to strains and difficulties experienced by the caregiver of mentally ill patient and include a range of psychological, emotional, social, physical, and financial problems.<sup>6</sup>

The prevalence of care giver burden for spouses is high.<sup>7</sup> In India overall care giver burden for spouse was 54%, majority had experienced severe to extremely severe burden.<sup>8</sup> In Nepal, married caregiver felt higher burden than unmarried.<sup>9</sup>

This study aims to assess the burden among spouses of psychiatric patients, the domains and factors associated with the burden.

## Methods

A quantitative, cross-sectional study was conducted in psychiatry outpatient department of Patan hospital, Patan Academy of Health Science (PAHS), Lagankhel, Lalitpur, Nepal. Non-probability purposive sampling technique was used. Sample size was calculated by using prevalence of caregiver's burden.<sup>10</sup>

The data collection instruments consisted of two parts.

Part I: Socio demographic Questionnaire -9 questions- 1) Type of diagnosis 2) Respondent relation to patient 3) Completed age 4) Educational status 5) Type of family 6) Occupation 7) Husband/wife income 8) Duration of caring 9) Permanent address and temporary address.

Part II: Burden Assessment schedule (BAS)<sup>11</sup> developed & standardized by step wise ethnographic exploration method in India. The inter-rater reliability and content validity are

high 0.80. Criterion validity was established by comparing with Family Burden Interview Schedule ranging between 0.71-0.82. Nepal and India share similar socio-cultural aspects. Forward and back translation of the instrument was done by utilizing the guideline Wild et al. 2005.<sup>12</sup> Reliability of the tool was maintained by Cronbach's coefficient (r) is 0.86.

The ethical clearance was obtained from Institutional Review committee of (IRC) of PAHS. Permission was taken from Hospital director, Nursing director, head of department of psychiatry department and OPD in-charge of Patan hospital.

Informed verbal and written consent was taken and face-to-face interview was conducted. Data analyzed using the Statistical Program for Social Science version 16. Inferential statistics (Chi-square test) was used to determine the association between variables. Level of significance was set at 0.05.

## Results

There were 160 respondents. Mean burden score was  $86.03 \pm 10.58$ , and 56 (35%) caregiver were in age group 40-49 years ( $41.01 \pm 10.30$ ). Eighty six (54%) of the spouses were husband; 54 (34%) respondents had basic level of education; 3 (2%) had master degree; 103 (65%) living in nuclear family; 29 (19%) had job and 61 (39%) had other occupation (driver, tailor, singer etc.), 59 (39%) had income of NPR 10,000 to 20,000 per month and the mean duration of caring was  $3.78 \pm 5.04$  years, 75 (47%) had duration of caring of 6-11 months. Diagnosis of 44 (28%) of respondent's spouse was neurotic, stress-related & somatoform disorders.

In 94 (58.75%) the level of care giver burden was severe, Table 1; highest maximum score of 10 was obtained in the domain of physical and mental health, Table 2. The care giver burden was significantly associated with relation to patient, Table 3; income, Table 4; and diagnosis, Table 5.

**Table 1. Level of burden among spouse of psychiatric patient (n=160)**

Level of burden	Frequency	Percentage
Mild burden (41-60)	-	-
Moderate burden (61-80)	53	33.13
Severe burden (81-100)	94	58.75
Very severe burden (101-120)	13	8.13
Mean score $\pm$ SD=86.03 $\pm$ 10.58		

**Table 2. Mean score & Standard deviation of each domain of Burden Assessment Schedule (n=160)**

Domains	Total no. of items in the domain	Minimum score of each domain	Maximum score of each domain	Mean	SD
Spouse related	5	6.00	13.00	9.13	1.50
Physical & mental health	6	6.00	18.00	11.34	3.35
External support	5	5.00	15.00	9.49	2.39
Caregiver's routines	4	5.00	11.00	8.14	1.27
Support of pt.	3	3.00	9.00	6.40	1.10
Taking responsibility	4	5.00	12.00	9.80	1.21
Other relations	3	3.00	9.00	4.84	1.72
Patient's behaviour	4	4.00	12.00	7.88	2.41
Caregiver's strategy	4	5.00	12.00	8.53	1.36

SD: Standard Deviation

**Table 3. Association between Sociodemographic Characteristics (Age, Respondent relation to patient, Educational status, Type of family) of Spouse of psychiatric patient and level of burden (n=160)**

Variables	Moderate burden	Severe & very severe burden	Chi-square (x <sup>2</sup> )	p-value
	n (%)	n (%)		
Age (in year)				
19-39	25 (36.23%)	44 (63.77%)	5.29	0.46
40-79	28 (30.77%)	63 (69.23%)		
Respondents relation to patient				
Husband	41 (47.67%)	45 (52.33%)	17.76	0.00*
Wife	12 (16.22%)	62 (83.78%)		
Educational status				
Basic level	27 (27.27%)	72 (72.73%)	7.88	0.01*
Secondary level	16 (35.56%)	29 (64.44%)		
Higher level	10 (62.50%)	6 (37.50%)		
Type of family				
Single	31 (30.10%)	72 (69.90%)	1.19	0.27
Joint	22 (38.60%)	34 (61.40%)		

P value &lt;.05=significant

\*significant

**Table 4. Association between Sociodemographic Characteristics (Type of occupation, Husband/Wife income per month, Permanent address and Temporary address) of Spouse of psychiatric patient and level of burden, n=160**

Variables	Moderate burden n (%)	Severe & very severe burden n (%)	Chi-square (x <sup>2</sup> )	p-value
Type of occupation				
Agriculture	24 (32.88%)	49 (67.12%)	0.04	0.95
Job	29 (33.33%)	58 (66.67%)		
Husband/wife's income/month (n=152)				
<20 thousands	27 (27.00%)	73 (73.00%)	4.60	0.03*
>20 thousands	23 (44.13%)	29 (55.77%)		
Permanent address				
Municipality	46 (33.82%)	90 (66.18%)	0.20	0.65
Metropolitan	7 (29.17%)	17 (70.83%)		
Temporary address(n=70)				
Municipality	12 (38.71%)	19 (61.29%)	0.21	0.64
Metropolitan	13 (33.33%)	26 (66.67%)		

P value <.05=significant \*significant

**Table 5. Association between Caregiving information and Clinical Characteristics of spouse (duration of caring and Diagnostic category) of Spouse of psychiatric patient and level of burden (n=160)**

Variables	Moderate burden n (%)	Severe & very severe burden n (%)	Chi-square x <sup>2</sup>	p-value
Duration of caring				
<5 year	44 (35.77%)	79 (64.23%)	1.68	0.19
>5 year	9 (24.32%)	28 (75.68%)		
Diagnostic category				
Substance use disorder	4 (10.26%)	35 (89.74%)	14.73	0.05*
Schizophrenia, schizotypal & delusional disorder	6 (27.27%)	16 (72.73%)		
Mood disorders	17 (43.59%)	22 (56.41%)		
Neurotic stress-related & Somatoform disorders	18 (40.91%)	26 (59.09%)		
Others*	8 (50.00%)	8 (50.00%)		

Others\*Organic mental disorder, migraine headache, epilepsy P value <.05=significant \*significant

## Discussions

In this study mean burden score was 86.03±10.58, less than the findings of study conducted in India<sup>13</sup> in 2018 with mean score of 73.5±14.0 and study from Nepal<sup>9</sup> in 2015 with score of 71.78±6.56. Similarly, another study conducted in 2015 in Iran has contradictory findings that revealed that the mean average score to be 51.73±18.23.<sup>14</sup>

In this study, spouse felt highest burden on the domains like physical and mental health (mean score of 11.34±3.35) followed by taking responsibility (score 9.80±1.21) but lowest burden on relations (score 4.84±1.72). The study from India reports similar findings with maximum burden in the domains of physical

and mental health (7.48±1.74) followed by external support (6.81±1.60), caregiver routine (5.62±1.02), taking responsibility (5.43±1.53).<sup>15</sup> Another study from India also supports present study findings that the highest amount of burden was in areas of physical and mental health (11.98±3.27), spouse related (9.90±1.86), and external support (9.40±2.57).<sup>16</sup>

In this study, significant association was seen between severe and very severe burden score and wife as a care giver, having basic level of education, income of less than NPR 20,000 and patient diagnosed as substance use disorder. Similar findings were reported from study conducted in India where female spouses reported slightly higher level of burden than

male, those from lower socioeconomic status, those from rural areas but no significant association to the type of family.<sup>13</sup>

Similarly, a study conducted in Nepal in 2015 shows that no significant difference between level of burden and educational status, socioeconomic status, duration of care and type of illness.<sup>9</sup> Another study conducted in India in 2016 showed that clinical diagnosis did not elicit any statistically significant difference indicating no difference on burden experienced by the spouses based on the clinical diagnosis.<sup>17</sup> Another study conducted in India in 2014 had contradictory findings to present study that there was association between area of living and level of burden.<sup>18</sup>

Limitations of present study includes single center tertiary care university teaching hospital in urban Kathmandu valley, the capital city of Nepal, and may not represent the population in general.

## Conclusions

More than half of spouse of psychiatric patients had severe level of care giver burden. Female spouse, those who were less educated and income less than twenty thousand per month had higher burden.

## Acknowledgements

This research was done in partial fulfillment of my thesis for master in nursing, and I am thankful to my guide and supervisor for their suggestions and advices.

## Fundings

The study was funded by University Grant Commission, Nepal.

## Conflict of interests

None

## References

1. Vigo D, Thornicroft G, Atun R. Estimating the true global burden of mental illness. *Lancet Psychiatry*. 2016;3(2):171-8. DOI PubMed [GoogleScholar](#)
2. Vos T, Barber RM, Bell B, Bertozzi-Villa A, Biryukov S, Bolliger I, Charlson F, Davis A, Degenhardt L, Dicker D. Global Burden of Disease Study 2013 Collaborators. Global, regional, and national incidence, prevalence, and years lived with disability for 301 acute and chronic diseases and injuries in 188 countries, 1990-2013: a systematic analysis for the Global Burden of Disease Study 2013. *Lancet*. 2015;386(9995):743-800. DOI PubMed [GoogleScholar](#)
3. Swaroop N, Shilpa Ravi, Goud BR, Maria A, Pius TM, Pal A, et al. Burden among caregivers of mentally-ill patients: a rural community based study. *Int J Res Dev Health*. 2013;1(2):29-34. [GoogleScholar](#) [Weblink](#)
4. Upadhaya, KD. Mental health manual for health worker. Kathmandu: Center for Mental Health and Counseling; 2010.
5. Kumari S, Singh AR, Verma AN, Verma PK, Chaudhury S. Subjective burden on spouses of schizophrenia patients. *Ind Psychiatry J*. 2009;18(2):97-100. DOI PubMed [GoogleScholar](#)
6. Magliano L, Fiorillo A, De Rosa C, Malangone C, Maj M, National Mental Health Project Working Group. Family burden in long-term diseases: a comparative study in schizophrenia vs. physical disorders. *Soc Sci Med*. 2005;61(2):313-22. DOI PubMed [GoogleScholar](#)
7. Viana MC, Gruber MJ, Shahly V, Alhamzawi A, Alonso J, Andrade LH, et al. Family burden related to mental and physical disorders in the world: results from the WHO World Mental Health (WMH) surveys. *Braz J Psychiatry*. 2013;35(2):115-25. DOI PubMed [GoogleScholar](#)
8. Stanley S, Bhuvanewari GM, Bhakyalakshmi S. Mental health status and perceived burden in caregiving spouses of persons with psychotic illness (a study from India). *Social Work in Mental Health*. 2016;14(5):530-44. DOI [GoogleScholar](#)
9. Bhandari AR, Marahatta K, Rana M, Ojha SP, Regmi MP. Caregiving burden among family members of people with mental illness. *Journal of Psychiatrists' Association of Nepal*. 2015;4(1):36-42. DOI [GoogleScholar](#) [Weblink](#)

10. World Economic Forum, Harvard School of Public Health. Methodological appendix: The global economic burden of non-communicable diseases. Geneva: World Economic Forum, 2011. [Weblink](#)
11. Zahid MA, Ohaeri JU. Relationship of family caregiver burden with quality of care and psychopathology in a sample of Arab subjects with schizophrenia. *BMC Psychiatry*. 2010;10(1):71. [DOI PubMed](#) [GoogleScholar](#)
12. Thara R, Padmavati R, Kumar S, Srinivasan L. Instrument to assess burden on caregivers of chronic mentally ill. *Indian J Psychiatry*. 1998;40(1):21-9. [PubMed](#) [GoogleScholar](#)
13. Wild D, Grove A, Martin M, Eremenco S, McElroy S, Verjee-Lorenz A, Erikson P. Principles of good practice for the translation and cultural adaptation process for patient-reported outcomes (PRO) measures: report of the ISPOR Task Force for Translation and Cultural Adaptation. *Value in Health*. 2005;8(2):94-104. [DOI PubMed](#) [GoogleScholar](#) [Weblink](#)
14. Arun R, Inbakamal S, Tharyan A, Premkumar PS. Spousal caregiver burden and its relation with disability in schizophrenia. *Indian Journal of Psychological Medicine*. 2018;40(1):22-8. [DOI PubMed](#) [GoogleScholar](#)
15. Arciszewska AI, Siwek M, Dudek D. Caregiving burden and psychological distress among spouses of bipolar patients: comparative analysis of subtype I and II. *Psychiatria Polska*. 2015;49(6):1289-302. [DOI PubMed](#) [GoogleScholar](#) [Weblink](#) [Article in English, Polish]
16. Agarwal M, Kushwaha A, Nischal A, Kar SK, Gupta B, Tripathi A, Agarwal A. Burden of care in the caregivers of patients with anxiety disorders. *Indian Journal of Social Psychiatry*. 2017;33(1):22-8. [DOI Google Scholar](#)
17. Kumar S, Mohanty S. Spousal burden of care in schizophrenia. *Journal of the Indian Academy of Applied Psychology*. 2007;33(2):189-94. [GoogleScholar](#) [Weblink](#)