



ISSN: 2091-2749 (Print)  
2091-2757 (Online)

#### Correspondence

Dr. Nishchal Bogati  
Patan Hospital  
Patan Academy of Health  
Sciences, Lalitpur, Nepal  
Email:  
nishchalbogati@pahs.edu.np

#### Peer Reviewers

Prof. Dr Jay N Shah  
Patan Academy of Health  
Sciences

Asst. Prof. Dr Ashis Shrestha  
Patan Academy of Health  
Sciences

#### Submitted

30 Apr 2019

#### Accepted

22 Aug 2019

#### How to cite this article

Laxmi RC Karki, Nishchal Bogati.  
Age specific clinicopathological  
profile of ovarian mass. Journal  
of Patan Academy of Health  
Sciences. 2019Dec;6(2):18-22.

## Age specific clinicopathological profile of ovarian mass

Laxmi RC Karki<sup>1</sup>, Nishchal Bogati<sup>2</sup>

<sup>1</sup>Assoc. Prof., <sup>2</sup>Intern doctor, Dept. of Obstetrics and Gynecology, Patan Hospital, Patan Academy of Health Sciences, Lalitpur, Nepal

### Abstract

**Introductions:** Ovarian cancer is seventh leading cause of cancer death among women. Ovarian mass has age specific occurrence and may help in screening and management plans. This study was conducted to determine the age-wise clinical profile of the ovarian mass.

**Methods:** This review done to analyse the histologically diagnosed ovarian masses in cases operated during three years at Patan Hospital, Patan Academy of Health Sciences, Nepal. Clinical profile of the patients and age-wise distribution of histological types of ovarian mass were analyzed descriptively using Microsoft Excel.

**Results:** Peak incidence of ovarian tumor was in age group of 20-29 years with 93 (36%) cases out of 258 ovarian masses. All eight malignant tumors were in age group of  $\geq 40$  years. Neoplastic masses were 188 (72.9%), 95 (50.5%) germ cell tumor. In 70 (27.1%) non-neoplastic lesion, corpus luteal cyst were 24 (34.3%).

**Conclusions:** Peak incidence of ovarian tumor was seen in age group of 20-29 years. Germ cell tumors accounted for half of the neoplastic lesions. All malignant ovarian tumors were in found in age group  $\geq 40$  years.

**Keywords:** germ cell tumor, ovarian cancer, ovarian mass

## Introductions

Ovarian mass can occur in all age groups.<sup>1</sup> About 80% of all are benign and mostly seen in age group of 20-45 years.<sup>2</sup> Ovarian cancer accounts for 2.5% of all malignancies and is the 7<sup>th</sup> leading cause of cancer death among women worldwide.<sup>3</sup> There are 2 to 6 new cases per 100000 women per year in Asia.<sup>4</sup> Study from Nepal shows benign ovarian mass accounts for 80-90% with peak incidence in age group of 20-40 years.<sup>5-7</sup>

Many non-neoplastic lesions of the ovary mimic ovarian neoplasms clinically and radiologically, and therefore, a histological classification is important to decide appropriate therapy.<sup>8,9</sup> There is a clear age distribution for these masses, which can serve as a useful guide for appropriate screening and management strategies.

This study aims to analyse the age-wise distribution of the various types of ovarian masses in local setup to serve as a baseline data for clinicians to develop screening and management protocols.

## Methods

This was a cross-sectional study done at Department of Obstetrics and Gynecology, Patan Hospital, Patan Academy of Health Sciences (PAHS), Nepal, during three years

period from January 2015 to December 2017. Data were collected from the hospital database with discharge diagnosis of ovarian mass. The individual files for each patient were obtained from the hospital record section and only those with histopathology report were included for analysis. World Health Organization (WHO) classification of ovarian tumors 2002 was used for classifying the tumors.<sup>10</sup> Ethical approval was obtained from the institutional review committee (IRC) of PAHS. Descriptive analysis was done using Microsoft Excel.

## Results

Out of total 258 ovarian masses, 93 (36%) were in age group 20-29 years, followed by 61 (23.6%) in 30-39 years. All eight (3.1%) malignant tumors were seen in age group  $\geq$ 40 years and among them, five in age group of 50-59 years. Mature cystic teratoma were 93 (36% of 258 ovarian mass), all benign, Table 1.

Out 258 ovarian masses, non-neoplastic were 70 (27.1%), Corpus luteal cyst 24 (34.3 %), Table 2.

Among 188 (72.9) neoplastic lesions, 177 (94.1%) were benign, 8 (4.3%) malignant and 3 (1.6%) borderline, Table 3.

**Table 1. Age distribution of neoplastic ovarian mass (N=258)**

Age years	Ovarian mass, total 258					N (%)
	Non-neoplastic 70 (27.13%)	Neoplastic Tumor 188 (72.9%)				
		Germ Cell	Surface Epithelial	Sex-cord Stromal	Metastatic	
$\leq$ 19	8	12	5	0	0	25 (9.7 %)
20-29	28	43	21	1	0	93 (36 %)
30-39	14	20	27	0	0	61 (23.6 %)
40-49	20	11	18	0	0	49 (19 %)
50-59	0	6	12	1	1	20 (7.7 %)
60-69	0	3	5	0	1	9 (3.5 %)
$\geq$ 70	0	0	1	0	0	1 (0.4 %)

**Table 2. Distribution of various non-neoplastic tumors (N=70)**

Types of non-neoplastic lesions	N (%)
Corpus luteal cyst	24 (34.3 %)
Simple cyst	18 (25.7%)
Endometrioid cyst	13 (18.6 %)
Hemorrhagic cyst	12 (17.1%)
Follicular cyst	3 (4.3 %)

**Table 3. Distribution of various neoplastic ovarian tumors (N=188)**

Neoplastic ovarian tumors	Nature	Types of lesion	N (%)
Germ cell tumor, 95 (50.5%)	Benign	Mature cystic teratoma	93 (49.5%)
		Struma ovarii	2 (1%)
Surface epithelial tumor, 89 (47.3%)	Benign	Serous	52 (27.7%)
		Mucinous	26 (13.8%)
	Borderline	Serous	1 (0.5%)
		Mucinous	2 (1%)
	Malignant	Serous	5 (2.7%)
		Mucinous	3 (1.6%)
Sex-cord stromal tumor, 2 (1%)	Benign	Fibroma	2 (1%)
Metastatic tumor, 2 (1%)			2 (1%)

## Discussions

This study showed most common age group for occurrence of ovarian tumor was in 20-29 years (36%), followed 30-39 years (23.6%), similar with the findings of other studies.<sup>6,11,12,18,19</sup> Ovarian tumor may occur at any age, including infancy and childhood.<sup>18</sup> The peak incidence is between 21 to 40 years and about 80% are benign.<sup>2,12</sup> This study showed that in age group of  $\leq 19$  years 48% were germ cell tumors, less common than the 95% reported in a study 'germ cell teratoma in children'.<sup>20</sup>

All eight malignant tumors in this study were found in  $\geq 40$  years. Malignant tumors are reported more commonly in older women, between the ages of 45 and 65 years.<sup>2</sup> Ovarian tumors are challenging as both non-neoplastic and neoplastic lesions present with similar clinical and radiological features. It is one of the reasons for prophylactic oophorectomies and hysterectomies. Early diagnosis is difficult due to its asymptomatic nature, inaccessible site and the limited use of various new techniques like cytology and

biopsy.<sup>9</sup> More than 70% of women with ovarian cancer are diagnosed in advanced stage with high fatality rate. Five-year survival rates for women with advanced disease range from 20% to 30% and early stages the cure rates are as high as 70% to 90%, and better survival related with age at presentation.<sup>21</sup>

This study showed a peak incidence of benign ovarian tumor between 2<sup>nd</sup> and 4<sup>th</sup> decade of life and malignant tumor after 4<sup>th</sup> decade. Thus, screening test like CA-125, abdominal and transvaginal sonography seems more appropriate in this age group of above 40 years for early detection of malignant ovarian mass.

In this study, 72.9% cases were neoplastic, and among them only 4.3% were malignant, less than 15.7%<sup>7</sup> reported earlier from our hospital on 363 ovarian tumors during three years data 2011-2013. One possible explanation could be, in earlier study the pathology department used to process samples from other hospitals in and around Kathmandu valley. However, this study did not look in depth to analyze the detail

demography of the samples which may represent geographical variation of malignancy or only suspicious malignant samples from outside were sent to our center for final histological confirmation.

In present study majority of non-neoplastic tumors were corpus luteal cyst (34.3%) followed by simple cyst of ovary (25.7%). Non-neoplastic lesions are commonly occurring ovarian lesions.<sup>12</sup> Similar findings of corpus luteal cyst have been reported.<sup>13,15</sup>

Ovarian tumor has heterogeneity in histological pattern although looks similar in gross examination and WHO classification is

also based on histogenesis of normal ovary. Surface epithelial tumors are the commonest according to this classification comprising 48.8% and 63.5% of all ovarian tumors in different studies while their malignant counterparts comprise approximately 90% of all ovarian cancers.<sup>12</sup>

But in present study, majority of tumors according to the WHO classification were of germ cell origin (51.6 %) followed by surface epithelial tumor which correspond to the earlier findings from our hospital<sup>7</sup> but contradict to other studies<sup>5,6,12</sup> in which majority of tumors were of surface epithelial origin, Table 4.

**Table 4. Comparative incidence of neoplastic ovarian tumors**

Types of ovarian tumor	R Jha et al. <sup>5</sup>	SK Maondal et al. <sup>12</sup>	S Vaidya et al. <sup>7</sup>	Current Study
Surface Epithelial Tumor	52.2%	67.9%	43.5%	47.3%
Germ-cell Tumor	42.2%	23.1%	51.5%	51.6%
Sex-cord Stromal Tumor	3.1%	5.6%	3.3%	1.1%

The variation in types of tumor in different studies from Nepal may be due to the differences in risk factors or exposures and demographic characteristics and demands further studies.

In present study our main focus was age group and ovarian mass, and analysis of further risk factors in multicentric pooled data may provide better generalizability.

## Conclusions

Our findings show benign ovarian tumors were more common than malignant, and germ cell tumors were common histologic type in all age groups. Peak incidence of ovarian tumor was in age group of 20-29 years. Mature cystic teratoma was the most common benign tumor. Serous carcinoma was the commonest malignancy occurring after 4<sup>th</sup> decade of life peaking between age of 50-59 years.

## Conflict of interests

None

## Fundings

None

## References

1. Rathod GB, Jain A. Non-neoplastic lesions of the ovary – a review. *International Archives of Integrated Medicine*. 2015;2(8):124-9. [GoogleScholar](#) [Weblink](#)
2. Kumar V, Abbas AK, Aster JC. Robbins and Cotran Pathologic basis of disease. 9th ed. Philadelphia, Pa: Elsevier Saunders; 2015. Chapter 22, The Female Genital Tract; p. 1022-34.
3. Reid BM, Permuth JB, Sellers TA. Epidemiology of ovarian cancer: a review. *Cancer Biol Med*. 2017;14(1):9-32. DOI [PubMed](#) [GoogleScholar](#)
4. Murad A. Ovulation induction and ovarian tumor: the debate continues. *Journal of the Pakistan Medical Association*. 1998;48(11):353-6. [PubMed](#) [GoogleScholar](#) [Weblink](#)

5. Jha R, Karki S. Histological pattern of ovarian tumors and their age distribution. *Nepal Med Coll J.* 2008;10(2):81-5. [GoogleScholar](#) [Weblink](#)
6. Kayastha S. Study of ovarian tumours in Nepal Medical College Teaching Hospital. *Nepal Med Coll J.* 2009;11(3):200-2. [GoogleScholar](#) [Weblink](#)
7. Vaidya S, Sharma P, KC S, Vaidya SA. Spectrum of ovarian tumors in a referral hospital in Nepal. *Journal of Pathology of Nepal.* 2014;4:539-43. [DOI](#) [GoogleScholar](#) [Weblink](#)
8. Gupta N, Bisht D, Agarwal AK, Sharma VK. Retrospective and prospective study of ovarian tumours and tumour-like lesions. *Indian J Pathol Microbiol.* 2007;50(3):525-7. [DOI](#) [PubMed](#) [GoogleScholar](#) [Weblink](#)
9. Kanthikar SN, Dravid NV, Deore PN, Nikumbh DB, Suryawanshi KH. Clinico-histopathological analysis of neoplastic and non-neoplastic lesions of the ovary: a 3-year prospective study in Dhule, North Maharashtra, India. *J Clin Diagn Res.* 2014;8(8):FC04-7. [PubMed](#) [Weblink](#)
10. FA Tavassoli, P Devillee. *World Health Organization Classification of Tumors. Pathology and Genetics of Tumors of Breast and Female Genital Organs.* Lyon: IARC Press; 2003. p. 113-96.
11. Pudasaini S, Lakhey M, Hirachand S, Akhter J, Thapa B. A study of ovarian cyst in a tertiary hospital of Kathmandu valley. *Nepal Med Coll J.* 2011;13(1):39-41. [GoogleScholar](#) [Weblink](#)
12. Mondal SK, Banyopadhyay R, Nag DR, Roychowdhury S, Mondal PK, Sinha SK. Histologic pattern, bilaterality and clinical evaluation of 957 ovarian neoplasms: a 10-year study in a tertiary hospital of eastern India. *J Cancer Res Ther.* 2011;7(4):433-7. [DOI](#) [PubMed](#) [GoogleScholar](#)
13. Shadab S, Tadayon T. Histopathological diagnosis of ovarian mass. *Journal of Pathology of Nepal.* 2018;8(1):1261-4. [DOI](#) [GoogleScholar](#) [Weblink](#)
14. Sawant A, Mahajan S. Histopathological study of ovarian lesions at a tertiary health care institute. *MVP Journal of Medical Sciences.* 2017;4(1):26-9. [DOI](#) [GoogleScholar](#) [Weblink](#)
15. Amatya S, Gurung G, Rana A. Annual clinicopathological analysis of ovarian tumors at TUTH. *NJOG.* 2010;4(2):18-24. [PDF](#) [Weblink](#)
16. NA Shaikh, Hashmi F, Samoo RP. Pattern of ovarian tumors: report of 15 years experience at Liaquat University Jamshoro. *J Liaquat Uni Med Health Sci.* 2007;6:13-5. [GoogleScholar](#) [Weblink](#)
17. Pradhan A, Sinha AK, Upreti D. Histopathological patterns of ovarian tumors at BPKIHS. *Health Renaissance.* 2012;10(2):87-97. [DOI](#) [Weblink](#)
18. Jindal D, Sahasrabhojane M, Jindal M, D'Souza J. Epidemiology of epithelial ovarian cancer: a tertiary hospital-based study in Goa, India. *Int J Reprod Contracept Obstet Gynecol.* 2017;6(6):2541-6. [DOI](#) [Weblink](#)
19. Maurya G, Singh SK, Pandey P, Chaturvedi V. Pattern of neoplastic and non-neoplastic lesions of ovary: a five-year study in a tertiary care centre of rural India. *Int J Res Med Sci.* 2018;6(7):2418-22. [DOI](#) [GoogleScholar](#) [Weblink](#)
20. Łuczak J, Bałaj M. Ovarian teratoma in children: a plea for collaborative clinical study. *J Ovarian Res.* 2018;11(1):75. [DOI](#) [PubMed](#) [GoogleScholar](#)
21. Rauh-Hain JA, Krivak TC, del Carmen MG, Olawaiye AB. Ovarian cancer screening and early detection in the general population. *Rev Obstet Gynecol.* 2011;4(1):15-21. [PubMed](#) [GoogleScholar](#)